

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



M

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶ : A61K 38/17, 38/43, C07H 21/04, C07K 14/435, 14/705, 16/28, 16/40, G01N 33/53, 33/543, 33/554	A1	(11) International Publication Number: WO 99/59620 (43) International Publication Date: 25 November 1999 (25.11.99)
(21) International Application Number: PCT/US99/10957 (22) International Filing Date: 19 May 1999 (19.05.99) (30) Priority Data: 60/086,155 19 May 1998 (19.05.98) US 60/124,070 12 March 1999 (12.03.99) US (71) Applicant: DUKE UNIVERSITY [US/US]; 230 North Building, Research Drive, Box 90083, Durham, NC 27708-0083 (US). (71)(72) Applicant and Inventor: STACK, Mary, S. [US/US]; 5434 North Lamon Avenue, Chicago, IL 60630 (US). (72) Inventors: MOSER, Tammy, L.; Duke University, 230 North Building, Research Drive, Box 90083, Durham, NC 27708-0083 (US). PIZZO, Salvatore, V.; Duke University, 230 North Building, Research Drive, Box 90083, Durham, NC 27708-0083 (US). (74) Agent: WILSON, Mary, J.; Nixon & Vanderhye P.C., 8th floor, 1100 North Glebe Road, Arlington, VA 22201-4714 (US).	(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: ANGIOSTATIN RECEPTOR (57) Abstract <p>The present invention relates, in general, to an angiotatin receptor and, in particular, to an angiotatin receptor present on cellular plasma membranes. More particularly, the present invention relates to the human angiotatin receptor, ATP synthase, or subunit or portion thereof, and to the use thereof in assays designed to screen compounds for their ability to serve as agonists or antagonists of human angiotatin. The invention further relates to nucleic acid sequences encoding ATP synthase, or subunit or portion thereof, and to host cells transformed therewith. The invention also relates to antibodies specific for ATP synthase.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

ANGIOSTATIN RECEPTOR

The present application claims priority from Provisional Application No. 60/086,155, filed May 19, 1998, and Provisional Application No. 60/124,070, filed March 12, 1999, the entire contents of both applications being incorporated herein by reference.

TECHNICAL FIELD

The present invention relates, in general, to an angiostatin receptor and, in particular, to an angiostatin receptor present on cellular plasma membranes. More particularly, the present invention relates to the human angiostatin receptor, ATP synthase, or subunit or portion thereof, and to the use thereof in assays designed to screen compounds for their ability to serve as agonists or antagonists of human angiostatin. The invention further relates to nucleic acid sequences encoding ATP synthase, or subunit or portion thereof, and to host cells transformed therewith. The invention also relates to antibodies specific for ATP synthase.

BACKGROUND

Tumor growth requires persistent blood vessel generation in the process of angiogenesis. If vascularization is prevented, tumor growth is dramatically impaired and the tumor size is restricted. Modulation of endogenous angiogenic inhibitors thus plays an important

role in tumor development. Angiostatin, a proteolytic fragment of plasminogen, is a potent inhibitor of angiogenesis and the growth of tumor cell metastases (O'Reilly et al, Cell 79:315-328 (1994)). Angiostatin
5 can be generated *in vitro* by limited proteolysis of plasminogen (Sottrup-Jensen et al, Progress in Chemical Fibrinolysis and Thrombolysis 3:191-209 (1978)) resulting in a 38kDa plasminogen fragment (Val⁷⁹-Pro³⁵³). Although the enzymatic mechanism by which angiostatin is generated
10 *in vivo* is unknown, recent studies have demonstrated that the cleavage of plasminogen to angiostatin can be catalyzed by a serine proteinase (Gately et al, Cancer Research 36:4887-4890 (1996)) and a macrophage metalloelastase (Dong et al, Cell 88:801-810 (1997)).
15 Generation of angiostatin from reduction of plasmin has also been shown *in vitro* (Gately et al, PNAS 94:10868-10872 (1997)) and in Chinese hamster ovary and human fibrosarcoma cells (Stathakis et al, JBC 272(33):20641-20645 (1997)).
20 Cellular receptors for plasminogen have previously been demonstrated on human umbilical vein endothelial cells (HUVEC) and are believed to function in the regulation of endothelial cell activities, including angiogenesis (Hajjar et al, J. Biol. Chem. 261(25):11656-
25 11662 (1986), Hajjar et al, JBC 269(3):21191-21197 (1994)). Receptors for plasminogen are also expressed in high numbers on tumor cells, where they have been identified as critical for tumor invasion. Proteins normally found in the cytoplasm have also been shown on

cell surface membranes and serve as plasminogen binding sites (Miles et al, Biochemistry 30:1682-1691 (1991)).

The present invention results from the demonstrations that plasminogen and angiostatin bind to distinct sites on cellular plasma membranes and that ATP synthase is the angiostatin binding protein. These findings make possible assays that can be used to screen compounds for their ability to modulate angiostatin activities. Compounds so identified have profound usefulness as therapeutic agents.

SUMMARY OF THE INVENTION

The present invention relates to an angiostatin receptor present on cellular plasma membranes. More particularly, the present invention relates to the human angiostatin receptor, ATP synthase, and to the use thereof, or of subunits or portions thereof, in assays designed to screen compounds for their ability to modulate angiostatin activities. The invention further relates to a nucleic acid sequences encoding ATP synthase, or subunit or portion thereof, and to host cells transformed therewith. The invention also relates to antibodies specific for ATP synthase.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A and 1B. Direct binding assay and Scatchard analysis of plasminogen and angiostatin with endothelial cells. HUVEC were plated at a density of

10,000 cells/well and incubated with increasing concentrations of ^{125}I -labeled-plasminogen or angiotatin. Fig. 1A. ^{125}I -labeled plasminogen binding was concentration-dependent and saturable with an apparent dissociation constant (K_d) of 158 nM and 870,000 sites/cell. Fig. 1B. Binding to HUVEC with ^{125}I -labeled angiotatin was concentration-dependent and saturable with a K_d of 245 nM and 38,000 sites/cell. Error bars represent standard deviation.

Figure 2. Competition binding assay between plasminogen and angiotatin. HUVEC were plated at a density of 10,000 cells/well and incubated with 1.0 μM ^{125}I -labeled plasminogen in the presence of 100-fold molar excess of unlabeled angiotatin for 1 h at 4°C. Cells were washed and the remaining radioactivity was quantified by γ -counting. (A) Total binding of 1.0 μM ^{125}I -labeled plasminogen was designated as 100%. (B) Plasminogen binding is inhibited by ~80% in the presence of a 25-fold molar excess of unlabeled plasminogen. (C) Plasminogen binding was not inhibited in the presence of a 100-fold molar excess of unlabeled angiotatin suggesting distinct binding sites for each on the cells. Similar experiments using ^{125}I -labeled angiotatin (D) showed no inhibition of binding in the presence of a 2-fold molar excess unlabeled plasminogen (E). Error bars represent standard deviation.

WO 99/59620

Figures 3A-3D. Affinity purification of plasminogen and angiostatin binding sites. SDS-PAGE containing membrane proteins were prepared and then analyzed by Western blotting. Membranes were incubated in 10 mM Tris-HCl, 0.15M NaCl, 0.05% NP40, pH 7.5 containing, Fig. 3A, streptavidin-alkaline phosphatase conjugate antibody, or, Fig. 3B, anti-annexin II antibody and developed using 5-bromo-4-chloroindol-3-yl-phosphate nitro blue tetrazolium. Membrane stained with Coomassie Brilliant blue, Fig. 3C, showing affinity purified membrane proteins. Membrane incubated with ^{125}I -labeled plasminogen, Fig. 3D, showing binding to the plasminogen purified membrane and not the angiostatin. Lane 1 represents protein eluted from the plasminogen-Sepharose column. Lane 2 represents protein eluted from the angiostatin-Sepharose column. The relative molecular weights of α -ATP synthase and β -ATP synthase are ~55 and ~50 kDa, respectively.

Figures 4A-4D. Binding of antibody directed against the α subunit of ATP synthase on the surface of HUVEC by flow cytometry. HUVEC were analyzed by FACScan Flow Cytometry. Histogram plots are shown for HUVEC (Fig. 4A) and A549 (Fig. 4B) where (--) represents cells incubated with antibody directed against the α subunit of ATP synthase, (---) pre-immune serum and (-) secondary antibody only. Histogram plot of A549 shown in Fig. 4C are similar with (--)

representing antibody incubated with a 5-fold molar excess α ATP synthase protein. In Fig. 4D, HUVEC demonstrate specific, saturable binding of antibodies directed against the α subunit of ATP synthase. The mean relative fluorescence of HUVEC incubated with pre-immune rabbit serum subtracted from the mean relative fluorescence of HUVEC incubated with the same volume of anti- α ATP synthase gave the mean relative fluorescence resulting from the specific binding of antibodies directed against the α subunit of ATP synthase on the HUVEC surface.

Figures 5A-5F. Immunofluorescence microscopy of ATP-synthase on HUVEC surface. HUVEC were incubated with rabbit polyclonal anti-serum raised against the α subunit of ATP synthase from *E. coli*. Fig. 5A, HUVEC under epi-illumination showing immunofluorescent surface staining for the α subunit of ATP synthase. Fig. 5B, Same field of HUVEC under visible light. Fig. 5C, Human dermal microvascular endothelial cells also showed immunofluorescent surface staining for the α subunit of ATP synthase. Control experiments were performed with Fig. 5D, pre-immune serum and Fig. 5E, secondary antibody alone. Fig. 5F, HUVEC were permeabilized by acetone fixation prior to adding antibodies for the α subunit of ATP synthase.

Figure 6. Competition binding assay between angiotstatin and the antibody against the α subunit of ATP synthase from *E. coli*. HUVEC were plated at a constant density of 10,000 cells/well and incubated with 0.5 μM ^{125}I -labeled angiotstatin in the presence of 1:10 dilution of antibody against the α subunit of ATP synthase from *E. coli* for 1 h at 4°C. Cells were washed and remaining bound radioactivity was quantified by γ -counting. Non-specific binding was measured in the presence of excess unlabeled angiotstatin and was subtracted from total binding. (A) Total binding of 0.5 μM ^{125}I -labeled angiotstatin was designated as 100%. (B) Angiotstatin binding is inhibited by 59% in the presence of a 1:10 dilution of anti- α subunit ATP synthase antibody. Competition studies were also performed simultaneously using rabbit pre-immune serum to account for non-specific inhibition. Error bars represent standard deviation. A 1 tailed homoscedastic t test was used for statistical analysis; $p < 0.10$.

Figures 7A-7E. Angiotstatin binding to the recombinant α subunit of human ATP synthase. The α subunit of human ATP synthase was cloned and expressed in *E. coli* and purified using Qiagen's nickel-Sepharose protein purification system before dialyzing in PBS, pH 7.0. Recombinant protein was electrophoresed on 5-15% SDS-PAGE, electroblotted onto Immobilon™ membrane and incubated 18 h in 10 mM Tris-HCl, 0.15M NaCl, 0.05% NP40, pH 7.5 (TSN) containing ^{125}I -angiotstatin. For

competition studies unlabeled ligand was added 4 h prior to radiolabeled ligand. Blots were washed in TSN buffer containing 0.05% Tween80 and bound radioactivity was quantified on a Molecular Dynamics
5 PhosphorImagerTM. Fig. 7A. Coomassie stain of Immobilon membrane containing the α subunit of human ATP synthase. Fig. 7B. Binding of 0.5 μ M ¹²⁵I-labeled angiostatin. Fig. 7C. Binding of 0.5 μ M ¹²⁵I-labeled
10 angiostatin in the presence of a 250-fold molar excess of unlabeled angiostatin. Binding of angiostatin is inhibited by ~56%. Fig. 7D. Binding of 0.5 μ M ¹²⁵I-labeled angiostatin in the presence of a 2500-fold molar excess of unlabeled plasminogen. Binding of
15 μ M ¹²⁵I-labeled plasminogen to the α subunit of human ATP synthase. Plasminogen did not bind to the recombinant α subunit of ATP synthase, however, it did bind the annexin II control (as shown in Fig. 3).

Figure 8. Binding of antibody directed against
20 the β subunit of ATP synthase on the surface of HUVEC by flow cytometry. HUVEC were analyzed by FACScan Flow Cytometry as described above and in the Examples. Histogram plots are shown for HUVEC cells incubated
25 with antibody directed against the β subunit of ATP synthase.

DETAILED DESCRIPTION OF THE INVENTION

The present invention results from the demonstration that plasminogen and angiostatin bind to distinct sites on the surface of endothelial cells, annexin II and ATP synthase, respectively. The invention provides methods for identifying compounds that can be used to modulate the effects of angiostatin *in vivo*, including its inhibitory effects on angiogenesis and endothelial cell migration and proliferation.

10 In one embodiment, the present invention relates to methods of screening compounds for their ability to bind ATP synthase and thereby to function, potentially, as angiostatin agonists or antagonists. ATP synthase includes two principal domains, an asymmetric membrane-
15 spanning F_0 portion containing a proton channel and a soluble F_1 portion containing three catalytic sites that cooperate in synthetic reactions. The F_1 region includes subunits α , β , γ , δ and ϵ . (See Elston et al, Nature 391:510 (1998).) The entire ATP synthase molecule can be
20 used in the present assays or α subunit thereof can be used, for example, the α and/or β subunit, the angiostatin binding domain of ATP synthase can also be used, as can a fusion protein comprising the synthase, the subunit thereof or the angiostatinn binding domain
25 thereof. The Examples that follow indicate that the α and β subunits of ATP synthase are present on the plasma membrane of endothelial cells. Further, the Examples indicate that angiostatin binds the α subunit. The α and

β subunits present on cellular plasma membranes may be identical to those present on mitochondrial membranes or they may represent a truncated (e.g., N- or C-terminal truncated) form thereof. The binding assays of the invention include the use of any such truncated forms.

Binding assays of this embodiment invention include cell-free assays in which ATP synthase, or subunit thereof or angiostatin binding domain thereof (or fusion protein containing same), is incubated with a test compound (proteinaceous or non-proteinaceous) which, advantageously, bears a detectable label (e.g., a radioactive or fluorescent label). Following incubation, the ATP synthase, or subunit thereof or angiostatin binding domain thereof (or fusion protein), free or bound to test compound, can be separated from unbound test compound using any of a variety of techniques (for example, the ATP synthase (or subunit or binding domain of fusion protein) can be bound to a solid support (e.g., a plate or a column) and washed free of unbound test compound). The amount of test compound bound to ATP synthase, or subunit thereof or angiostatin binding domain thereof (or fusion protein), is then determined, for example, using a technique appropriate for detecting the label used (e.g., liquid scintillation counting and gamma counting in the case of a radiolabelled test compound or by fluorometric analysis).

Binding assays of this embodiment can also take the form of cell-free competition binding assays. In such an assay, ATP synthase, or subunit thereof or angiostatin binding domain thereof, or fusion protein containing

- same, is incubated with a compound known to interact with ATP synthase (e.g., angiostatin or ATP synthase binding portion thereof), which compound, advantageously, bears a detectable label (e.g., a radioactive or fluorescent label). A test compound (proteinaceous or non-proteinaceous) is added to the reaction and assayed for its ability to compete with the known (labeled) compound for binding to ATP synthase, or subunit thereof or angiostatin binding domain thereof (or fusion protein).
- 10 Free known (labeled) compound can be separated from bound known compound, and the amount of bound known compound determined to assess the ability of the test compound to compete. This assay can be formatted so as to facilitate screening of large numbers of test compounds by linking
- 15 the ATP synthase, or subunit thereof or angiostatin binding domain thereof (or fusion protein), to a solid support so that it can be readily washed free of unbound reactants. A plastic support, for example, a plastic plate (e.g., a 96 well dish), is preferred.
- 20 ATP synthase, or subunit thereof or angiostatin binding domain thereof (or fusion protein), suitable for use in the cell-free assays described above can be isolated from natural sources (e.g., membrane preparations) or prepared recombinantly or chemically.
- 25 The ATP synthase, or subunit thereof or angiostatin binding domain thereof, can be prepared as a fusion protein using, for example, known recombinant techniques. Preferred fusion proteins include a GST (glutathione-S-transferase) moiety, a GFP (green fluorescent protein)
- 30 moiety (useful for cellular localization studies) or a

His tag (useful for affinity purification). The non-ATP synthase moiety can be present in the fusion protein N-terminal or C-terminal to the ATP synthase, subunit or binding domain.

5 As indicated above, the ATP synthase, or subunit thereof or angiostatin binding domain thereof, or fusion protein, can be present linked to a solid support, including a plastic or glass plate or bead, a chromatographic resin (e.g., Sepharose), a filter or a
10 membrane. Methods of attachment of proteins to such supports are well known in the art and include direct chemical attachment and attachment via a binding pair (e.g., biotin and avidin or biotin and streptavidin). It will also be appreciated that, whether free or bound to a
15 solid support, the ATP synthase, or subunit thereof or angiostatin binding domain thereof, or fusion protein, can be unlabeled or can bear a detectable label (e.g., a fluorescent or radioactive label).

The binding assays of the invention also include
20 cell-based assays in which ATP synthase, or subunit thereof or angiostatin binding domain thereof or fusion protein, is presented on a cell surface. Cells suitable for use in such assays include cells that naturally express ATP synthase and cells that have been engineered
25 to express ATP synthase (or subunit thereof or angiostatin binding domain thereof or fusion protein comprising same). The cells can be normal or tumorigenic. Advantageously, cells expressing human ATP synthase are used. Examples of suitable cells include
30 procaryotic cells (e.g., bacterial cells (e.g., *E.*

coli)), lower eucaryotic cells (e.g., yeast cells (e.g., hybrid kits from Promega (CG 1945 and Y190), and the strains YPH500 and BJ5457)) and higher eucaryotic cells (e.g., insect cells and mammalian cells (e.g.,
5 endothelial cells, including bovine aortic endothelial cells (BAEC), bovine adrenal medulla endothelial cells (BAMEC), murine endothelial cells CP_γ-4-1, HUVEC or any human endothelial cell line, or cells such as human lung carcinoma cells (e.g., A549 cells))).

10 Cells can be engineered to express ATP synthase (advantageously, human ATP synthase, or subunit thereof or angiostatin binding domain thereof, or fusion protein that includes same) by introducing into a selected host an expression construct comprising a sequence encoding
15 ATP synthase, or subunit thereof or angiostatin binding domain thereof or fusion protein, operably linked to a promoter. A variety of vectors and promoters can be used. For example, pET-24a(+) (Novagen) containing a T7 promoter is suitable for use in bacteria, likewise,
20 pGEX-5X-1. Suitable yeast expression vectors include pYES2 (Invitron). Suitable baculovirus expression vectors include p2Bac (Invitron). Suitable mammalian expression vectors include pBK/CMV (Stratagene). Introduction of the construct into the host can be
25 effected using any of a variety of standard transfection/transformation protocols (see Molecular Biology, A Laboratory Manual, second edition, J. Sambrook, E.F. Fritsch and T. Maniatis, Cold Spring Harbor Press, 1989). Cells thus produced can be cultured

using established culture techniques suitable for the involved host. Culture conditions can be optimized to ensure expression of the ATP synthase (or subunit, binding domain or fusion protein) encoding sequence.

5 While for the cell-based binding assays the ATP synthase (or subunit, binding domain or fusion protein) can be expressed on a host cell membrane (e.g., on the surface of the host cell), for other purposes the encoding sequence can be selected so as to ensure that
10 the expression product is secreted into the culture medium.

The cell-based binding assays of the invention can be carried out by adding test compound (advantageously, bearing a detectable (e.g., radioactive or fluorescent)
15 label), to medium in which the ATP synthase (or subunit thereof or angiotensin binding domain thereof or fusion protein containing same) expressing cells are cultured, incubating the test compound with the cells under conditions favorable to binding and then removing unbound
20 test compound and determining the amount of test compound associated with the cells.

Identification of ATP synthase on a cell membrane (e.g., on the cell surface) can be made using techniques such as those in the Examples that follow (e.g., the cell
25 surface can be biotin labeled and the protein followed by a fluorescent tag). Membrane associated proteins (e.g., cell surface proteins) can also be analyzed on a Western blot and the bands subjected to mass spectroscopy analysis. For example, a fluorescently tagged antibody
30 can be used with permeabilized cells which cells can then

be probed with another fluorescently tagged protein. Each tag can be monitored at a different wavelength, for example, using a confocal microscope to demonstrate co-localization.

5 As in the case of the cell-free assays, the cell-based assays can also take the form of competitive assays wherein a compound known to bind ATP synthase (and preferably labelled with a detectable label) is incubated with the ATP synthase (or subunit thereof or angiostatin
10 binding domain thereof or fusion protein comprising same) expressing cells in the presence and absence of test compound. The affinity of a test compound for ATP synthase can be assessed by determining the amount of known compound associated with the cells incubated in the
15 presence of the test compound, as compared to the amount associated with the cells in the absence of the test compound.

It will be appreciated from a reading of this disclosure that the selectivity of a test compound for
20 cell surface ATP synthase, as compared to mitochondrial ATP synthase, can be easily assessed. Compounds which, by virtue of their physicochemical properties, cannot diffuse across cellular membranes (and that are not natural or artificial ligands for cell transporters) can
25 be considered selective for cell surface ATP synthase. For example, compounds that bind cell surface ATP synthase but are positively charged can thereby be prevented from diffusing across membranes.

A test compound identified in one or more of the
30 above-described assays as being capable of binding to ATP

synthase can, potentially, mimic or enhance the effects of angiostatin on angiogenesis, cellular migration, proliferation and pericellular proteolysis or, potentially, antagonize the effects of angiostatin, for example, by precluding angiostatin from binding its receptor. To determine the specific effect of any particular test compound selected on the basis of its ability to bind ATP synthase (or inhibit (competitively or non-competitively) angiostatin binding to ATP synthase), assays can be conducted to determine, for example, the effect of various concentrations of the selected test compound on activity, for example, cell (e.g., endothelial cell) proliferation, metabolism or cytosolic/cytoplasmic pH. (Assays can be conducted to determine the effect of test compounds on ATP synthase (and ATPase) activity using standard enzyme assay protocols.)

Cell proliferation can be monitored by measuring uptake of labeled (e.g., radioactively (e.g., ^3H , ^{51}C , ^{14}C), for example, fluorescently (e.g., CYQUANT (Molecular Probes)) or colorimetrically (e.g., βrdU (Boehringer Mannheim or MTS (Promega)), bases into cellular nucleic acids. Cytosolic/cytoplasmic pH determinations can be made with a digital imaging microscope using substrates such as BCECF (bis(carboxyethyl)-carbonyl fluorescein) (Molecular Probes, Inc.) A test compound that reduces or replaces the concentration of angiostatin required to inhibit cellular proliferation or lower intracellular pH can be expected to do so by acting as an angiostatin agonist. A test compound that enhances cellular

proliferation in the presence of angiostatin (or functional portion thereof or functional equivalent thereof) can be expected to do so by acting as an angiostatin antagonist. A test compound that raises intracellular pH in the presence of angiostatin (or functional portion thereof or functional equivalent thereof) may do so by acting as an angiostatin antagonist. These functional assays can also be conducted in the absence of angiostatin (i.e., test compound alone), with angiostatin (or functional portion thereof or functional equivalent thereof) run as a separate control. A test compound that, for example, modulates intracellular pH in the absence of angiostatin can be an angiostatin agonist or antagonist.

Other types of assays that can be carried out to determine the effect of a test compound on angiostatin binding to ATP synthase include the Lewis Lung Carcinoma assay (O'Reilly et al, Cell 79:315 (1994)) and extracellular migration assays (Boyden Chamber assay: Kleinman et al, Biochemistry 25:312 (1986) and Albini et al, Can. Res. 47:3239 (1987)).

Das et al (J. Exp. Med. 180:273 (1994)) have reported the presence of the β subunit of H^+ transporting ATP synthase on the plasma membrane of human tumor cell lines. The present demonstration of the α subunit of ATP synthase on plasma membranes, and the binding thereto of angiostatin, indicates that angiostatin may be directly involved in effecting cytolysis, for example, of tumor cells. The binding of angiostatin to its receptor may

result in the transport of protons across plasma membranes and into cells with the result being cytolysis by osmotic shock. Accordingly, the present invention includes within its scope methods of screening compounds for their ability to modulate the effect of angiostatin on proton pumping that results from the binding of angiostatin to its receptor. In one such assay, cells that express ATP synthase (or subunit (e.g., α or β) or portion thereof) are incubated with the test compound in the presence of angiostatin (or functional portion thereof or functional equivalent thereof) and the influx of protons into the cells determined and compared to the influx of protons observed in the absence of the test compound. Compounds that reduce the concentration of angiostatin (or functional portion thereof or functional equivalent thereof) necessary to effect a particular level of proton influx can be expected to do so by acting as a angiostatin agonist. Compounds that reduce the amount of angiostatin-induced proton pumping observed can be expected to do so by acting as an angiostatin antagonist. The amount of proton pumping can be determined using any of a variety of approaches, including using cells preloaded with a pH sensitive reporter (for example, BCECF can be used to measure pH (Misra et al, Biochem. J. 309:151 (1995)) and monitoring the effect of the test compound on the reporter. Alternatively, the effect of a test compound on proton pumping can be determined by monitoring cell lysis using, for example, a chromium 51 release assay (McManus et al,

Exper. Lung Res. 15:849 (1989); Zucker et al, Res. Comm.
Chem. Path. Pharm. 39:321 (1983)).

In addition to the various approaches described
above, assays can also be designed so as to be
5 monitorable colorometrically or using time-resolved
fluorescence.

In another embodiment, the invention relates to
compounds identified using the above-described assays as
being capable of binding to ATP synthase (and/or
10 inhibiting angiostatin from binding to ATP synthase
(competitively or non-competitively) and/or modulating the
angiostatin effects on cellular bioactivities and/or
modulating ATP synthase activity. Such compounds can
include novel small molecules (e.g., organic compounds
15 (for example, organic compounds less than 500 Daltons),
and novel polypeptides, oligonucleotides, as well as
novel natural products (preferably in isolated form)
(including alkyls, tannins, glycosides, lipids,
carbohydrates and the like). Compounds that mimic or
20 enhance angiostatin activities can be used to inhibit
angiogenesis, for example, in tumor bearing patients and
in patients suffering from vascular related retinopathies
(including diabetic) and Terigium. Other diseases in
which angiogenesis is a significant component of tissue
25 pathology include rheumatoid arthritis and keloid
formation. Compounds that inhibit angiostatin activities
can be used to promote angiogenesis in conditions of
vascular insufficiency, including ischemic heart disease,
peripheral vascular disease, thromboembolic disease,
30 stroke and vasculitities (Buerger's disease, Wegener's

granulomatosis, and Giant Cell Arteritis). Such compounds can also be used at wound sites to promote healing, and at sites of transplantation and grafting (e.g., skin grafting). Other diseases/disorders amenable to treatment using compounds selected in accordance with the above-described assays include obesity, osteoarthritis, vascular diseases/disorders of the eye, including diabetic retinopathy, macular degeneration, retinopathy of prematurity, corneal inflammation, and viral infections, as well as psoriasis, spinal cord injuries, and other diseases and disorders that can be expected to benefit from intervention of vascularization.

The compounds identified in accordance with the above assays can be formulated as pharmaceutical compositions. Such compositions comprise the compound and a pharmaceutically acceptable diluent or carrier. The compound can be present in dosage unit form (e.g., as a tablet or capsule) or as a solution, preferably sterile, particularly when administration by injection is anticipated. The compound can also be present as a cream, gel or ointment, for example, when topical administration is preferred. The dose and dosage regimen will vary, for example, with the patient, the compound and the effect sought. Optimum doses and regimens can be determined readily by one skilled in the art.

In a specific embodiment, the invention relates to a method of antagonizing the effect of angiostatin in a patient by administering ATP synthase, or soluble, angiostatin binding portion thereof. The ATP synthase, or portion thereof, suitable for use in this method can

be prepared recombinantly or chemically and can be formulated with an acceptable carrier (including a liposome) as a pharmaceutical composition. The ATP synthase, or portion thereof, can be present as a fusion protein, for example, fused to the heavy chain of IgG. The ATP synthase, or portion thereof, can be derivatized (e.g., with polyethylene glycol) so as to modify its half life in vivo. The method of this embodiment finds application, for example, in wound healing.

10 In another embodiment, the invention relates to antibodies specific for ATP synthase, and antigen binding fragments thereof, including F(ab)₂' or F(ab) fragments. The antibodies can be monoclonal or polyclonal and can be prepared using standard techniques (Harlow and Lane
15 Antibodies, A Laboratory Manual, (1988) Cold Spring Harbor Laboratories). The antibodies can be used in ATP purification protocols or the antibodies can be formulated as pharmaceutical compositions and used therapeutically to mimic or enhance the effects of
20 angiostatin on endothelial cells or to antagonize such effects.

In yet a further embodiment, the invention relates to kits, for example, kits suitable for conducting assays described herein. Such kits can include ATP synthase, or
25 subunit thereof or angiostatin binding domain thereof, or fusion protein comprising same, and/or angiostatin. These components can bear a detectable label. The kit can include an ATP synthase-specific or angiostatin-specific antibody. Plasminogen can also be present. The
30 kit can include any of the above components disposed

within one or more container means. The kit can further include ancillary reagents (e.g., buffers) for use in the assays.

5 In another embodiment, the present invention relates to methods of diagnosis that are based, for example, on assays for the binding of angiostatin to ATP synthase. Such methods make possible the identification of patients suffering from diseases, disorders or conditions associated with abnormal angiogenesis.

10 The demonstration that ATP synthase is the angiostatin binding protein, and the resulting availability of methods of identifying agents that can be used to modulate the effects of angiostatin, make it possible to determine which individuals will likely be
15 responsive to particular therapeutic strategies. Treatment strategies for individuals suffering from a disease, disorder or condition associated with abnormal angiogenesis can thus be designed more effectively and with greater predictability of a successful result.
20 Thus, for a given clinical disease that is associated both with abnormal angiogenesis and that is of polygenic (non-Mendelian) origin, one would select that genotype that is implicated not only in the disease, but also in that variant of the disease that is
25 associated with abnormal angiogenesis and proceed to screen, via a diagnostic procedure, all future patients having the same genotype in order to choose that therapeutic strategy most associated with a successful outcome or least associated with a toxic side effect,
30 for that genotype.

Certain aspects of the present invention are described in greater detail in the non-limiting Examples that follow.

EXAMPLES

5 The following experimental details are referenced in the specific Examples that follow.

Protein Purification

Plasminogen was purified from human plasma by affinity chromatography and separated into isoforms 1 and
10 2 as previously described (Deutsch et al, Science 170:1095 (1970), Gonzalez-Gronow et al, Biochemistry 23:190-194 (1984)). Based on kinetic and electrophoretic analysis, all plasminogen preparations were shown to be plasmin-free. The concentration of plasminogen was
15 determined spectrophotometrically at 280 nm using an $A^{1\%}_{1\text{cm}}$ value of 1.67 and a molecular mass of 92 kDa for Glu¹-plasminogen (Castellino et al, Chem. Rev. 81:431 (1981)). Human plasminogen kringle 1-3 (angiotatin) were purified as previously described (Sottrup-Jensen et
20 al, Progress in Chemical Fibrinolysis and Thrombolysis 3:191-209 (1978)). The concentration of angiotatin was determined spectrophotometrically at 280 nm using an $A^{1\%}_{1\text{cm}}$ value of 0.8 and a molecular mass of 38 kDa (Sottrup-Jensen et al, Progress in Chemical Fibrinolysis
25 and Thrombolysis 3:191-209 (1978)). Protein endotoxin levels were <50.0 pg endotoxin/ml as assessed by Pyrotell *Limulus* amoebocyte lysate clotting times (Associates of Cape Cod, Woods Hole, MA)

Cell Culture

Primary human umbilical vein endothelial cells (HUVEC) were grown as previously described (Morales et al, Circulation 91:755-763 (1995)) in 150 mm petri dishes and retained for up to 6 passages. Human dermal microvascular endothelial cells (HMVEC) were obtained from Clonetics (San Diego, CA), grown according to specifications and retained for up to 6 passages. A549 (human lung carcinoma) cells were obtained from ATCC (Rockville, MD) and grown according to specifications. For all experiments cells were detached by incubation with PBS containing 2 mM EDTA, pH 7.4.

Antibody Purification

Antibody to His-tagged recombinant α -subunit ATP synthase was generated in rabbits by intranodal injection (Covance Laboratories, Vienna, VA). Production bleeds were centrifuged and the serum obtained was ammonium sulfate precipitated. The precipitate was resuspended in PBS/0.5M NaCl, pH 7.5 and passed over Protein A-Sepharose (Sigma, St. Louis, MO), plasminogen-Sepharose and α -subunit ATP synthase-Sepharose columns (CNBr coupling, Pharmacia Amersham, Piscataway, NJ). Each column was eluted with 20 mM glycine, pH 2.5. Neutralized IgG fractions were tested by immunodiffusion, ELISA and Western blotting. Antibody to the α -subunit of ATP synthase showed no cross-reactivity with plasminogen or other proteins by Western blot analysis. Polyclonal antibody obtained from Dr. A.E. Senior (Rochester Medical Center,

Rochester, NY) directed against the α subunit of ATP synthase from *E. coli* was characterized by ELISA and Western blot analysis and showed no cross-reactivity with other proteins in the F_1 portion or *E. coli* cell membranes (Perlin et al, Archives of Biochem. and Biophys. 236(2):603-611 (1985), Rao et al, Archives of Biochem. and Biophys. 255(2):309-315 (1987)).

Binding Assays

Ligands were radioiodinated using Iodobeads™ (Pierce), repurified on L-lysine-Sepharose, eluted with 100 mM ϵ -aminocaproic acid (EACA) and dialyzed in PBS, pH 7.0, before use in binding assays. HUVEC were plated at a density of 5000 or 10,000 cells/well and incubated with increasing concentrations of ^{125}I -labeled ligand in media containing 1% bovine serum albumin (BSA) for 1 h at 4°C in 96-well plates. Wells were washed and remaining bound radioactivity was quantified using an LKB 1272 γ -radiation counter. Non-specific binding was measured in the presence of excess unlabeled ligand.

Membrane Purification

Plasma membrane extracts from NHS-biotin labeled HUVEC were prepared by 300 psi Parr bomb nitrogen cavitation and ultracentrifugation (Young et al, J. Biol. Chem. 270(3):999-1002 (1995)). Membrane extracts were incubated with plasminogen-Sepharose or angiostatin-Sepharose columns in an inhibitor cocktail buffer (Young et al, J. Biol. Chem. 270(3):999-1002

(1995)). Each Sepharose column was eluted with 50 mM Tris/100mM EACA, pH 7.5, 50 mM Tris/1 M NaCl, pH 7.5, 50 mM Tris/7% DMSO and 20 mM glycine, pH 2.5 to account for all types of binding. The glycine eluates were
5 dialyzed, lyophilized, electrophoresed on 5-15% gradient SDS-PAGE (Laemmli, *Nature (London)* 227:680-685 (1970)) and electroblotted onto Immobilon™ membrane (Matsudaira, *J. Biol. Chem.* 262:10035-10038 (1987)) prior to experiments to identify plasminogen and
10 angiostatin binding proteins.

Mass Spectrometer Analysis

Plasma membrane proteins were separated on SDS-PAGE gels and the bands of interest were excised from the gels and digested *in situ* with trypsin. A portion
15 (1/20) of each sample was analyzed by MALDI-MS and the obtained mass spectrometric peptide maps were used to identify the protein in the OWL Protein database release 29.6 (Mann et al, *Biol. Mass Spectrom.* 22:338-345 (1993), Pappin et al, *Curr. Biol.* 3:327-332
20 (1993)).

Flow Cytometry

HUVEC and A549 cells were resuspended in ice-cold staining buffer (HBSS, 1% BSA, 0.1% sodium azide) and incubated on ice for 30 min with either rabbit
25 polyclonal anti-serum raised against α subunit ATP synthase derived from *E. coli* or pre-immune rabbit serum. Cells were washed with ice-cold staining buffer and pelleted in a microfuge at 4°C. This wash was

repeated twice and the cells resuspended in ice-cold staining buffer prior to incubation on ice for 30 min in the dark with goat anti-rabbit IgG conjugated to fluorescein isothiocyanate (FITC). Following the final wash (as above), the cells were pelleted and fixed in 10% neutral buffered formalin at a density of 1×10^6 cells/ml. Control experiments were performed using antibody directed against the α subunit of ATP synthase which was preincubated with a 5-fold molar excess of recombinant α subunit ATP synthase protein. The mean relative fluorescence following excitation at a wavelength of 488nm was determined for each sample on a FACScan flow cytometer (Becton-Dickenson) and analyzed with CellQuest software (Becton-Dickenson).

15 Immunofluorescence Microscopy

HUVEC and HMVEC were plated at 5×10^5 cells/ml on glass cover slips and allowed to adhere overnight. Cells were incubated at 4°C for 1 h in PBS, pH7.0 containing 1%BSA with either rabbit polyclonal anti-serum raised against the α subunit of ATP synthase derived from *E. coli*, pre-immune rabbit serum, pre-immune IgG, or anti-rabbit IgG. Cells were washed and incubated at 4°C for 1 h in the dark with goat anti-rabbit IgG conjugated to indocarbocyanine (Cy3) before washing and fixing in 4% paraformaldehyde. Immunofluorescence microscopy was performed using an Olympus BX-60 microscope (Olympus Corp., Lake Success, NY).

Cloning of the α Subunit of ATP synthase

Poly A⁺ mRNA was isolated from HUVEC using Oligotex resin (Qiagen). RNA was reverse transcribed into single-stranded cDNA using AMV Reverse

- 5 Transcriptase (Boeh. Mann.). The α subunit of ATP synthase was PCR amplified using Expand High Fidelity PCR system (Boeh. Mann.). The 1.7 kb PCR product was purified from a 0.8% TAE (tris-acetate/EDTA) agarose gel using a QIAEX II gel extraction kit. Restriction
- 10 enzyme digests of the PCR fragment and vector pLE1 were carried out at 37°C for 1 h. Both digests were passed over Qiaquick purification columns, then ligated overnight at 16°C using T4 DNA ligase. Competent *E. coli* DH5 α (Gibco BRL) were transformed with the
- 15 ligation mixture, plated on 2xYT agarose plates and grown overnight at 37°C. Colonies were screened for the insert via restriction enzyme digest and DNA sequencing.

Purification of the α Subunit of ATP Synthase

- 20 Competent *E. coli* BL21DE3 were transformed with the pLE1 vector containing the α -subunit, plated on 2xYT agarose and grown overnight at 37°C. 20ml of 2xYT (YT=bacto-yeast tryptone) containing 50 μ g/ml kanamycin were inoculated with one colony and grown overnight at
- 25 37°C (200 rpm). A one liter culture (2xYT, 50 μ g/ml kanamycin) was inoculated with 20 ml of the noninduced overnight culture and grown at 37°C to an A of 0.6 at a wavelength of 600nm. Isopropylthio-b-D-galactosidase

(IPTG) was added to a final concentration of 1mM and grown for an additional 3 hours. Cells were harvested by centrifugation at 8000 rpm for 10 minutes and stored overnight at -20°C. Lysates were prepared under
5 denaturing conditions and batch purified using Qiagen Ni-NTA agarose (Qiagen). Resulting protein was dialyzed against PBS, pH 7.0 for use in all experiments.

Proliferation Assay

10 HUVEC were plated at a density of 5000 cells/well in media depleted of fetal calf serum overnight to allow the cells to become quiescent. Fresh media containing fetal calf serum was added to the wells along with angiostatin at a final concentration of 0.5,
15 0.75 and 1.0 μ M. In some experiments, antibody directed against the α subunit of ATP synthase derived from *E. coli* was also added at a dilution of 1:10. MTS/PMS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl-2H-tetrazolium,
20 inner salt/phenazine methosulfate)) solution was added after 24 hours and the absorbance of formazan was quantitated on a Thermomax™ plate reader at a wavelength of 490nm according to the manufacturer's specifications (Promega, Madison, WI). The absorbance
25 values used to calculate the percent proliferation of the cells ranged from 0.81 for untreated, 0.60 for treated and 0.47 for baseline quiescent cells.

EXAMPLE 1

Competition Binding Assay

To determine whether angiostatin blocks angiogenesis by competitive interaction with endothelial cell plasminogen receptors, the effects of angiostatin on the binding of plasminogen to endothelial cells were analyzed.

In control experiments, plasminogen bound to HUVEC in a concentration-dependent saturable manner with an apparent dissociation constant (K_d) of 158 nM and ~870,000 sites per cell (Fig. 1A), comparable to values previously reported (Hajjar et al, JBC 269(3):21191-21197 (1994)). Angiostatin also bound to HUVEC in a concentration-dependent saturable manner with a similar affinity (K_d of 245 nM), with ~38,000 sites per cell (Fig. 1B). Binding studies utilizing ^{125}I -labeled plasminogen and a 100-fold molar excess of unlabeled angiostatin demonstrated no inhibition of plasminogen binding (Fig. 2). Similar studies were performed using ^{125}I -labeled angiostatin. Excess unlabeled plasminogen had little or no effect on angiostatin binding (Fig. 2). In contrast to plasminogen, binding of ^{125}I -labeled angiostatin to HUVEC in the presence of 100 mM ϵ -aminocaproic acid (EACA) was only slightly inhibited, suggesting that binding of angiostatin to endothelial cells is not a lysine binding site dependant process. Together, these data indicate the presence of a distinct angiostatin binding site on HUVEC.

EXAMPLE 2Purification of the Angiostatin Binding Site
From Endothelial Cells

The cell surface proteins involved in binding of plasminogen or angiostatin to HUVEC were identified by subjecting NHS-biotin labeled HUVEC plasma membranes to affinity chromatography on plasminogen-Sepharose or angiostatin-Sepharose. Two distinct bands were identified on Western blot analysis using streptavidin-alkaline phosphatase conjugate (Fig. 3A) or by Coomassie Brilliant blue stain (Fig. 3C). A companion blot, probed with an antibody to the known plasminogen receptor, annexin II, demonstrated immunologic cross-reactivity with the 44 kDa membrane protein isolated from the plasminogen-Sepharose column (Fig. 3B, lane 1), but not with the 55 kDa protein isolated from the angiostatin-Sepharose column (Fig. 3B, lane 2). Ligand blot analysis of the affinity purified plasma membranes using ^{125}I -labeled plasminogen (Fig. 3D, lanes 1 and 2) demonstrated binding of plasminogen only to the 44 kDa protein and not to the 55 kDa specie. These data further evidence the fact that HUVEC contains an angiostatin binding site distinct from the plasminogen binding protein, annexin II.

EXAMPLE 3Peptide Mass Fingerprinting of the
Angiostatin Binding Site

To identify the unique angiostatin binding site
5 component, the affinity-purified proteins were analyzed
by amino-terminal sequencing, mass spectrometer analysis
and peptide mass fingerprinting. Both the 44 and 55 kDa
proteins were analyzed by reduced SDS-PAGE and digested
with trypsin *in situ* (Matsui et al, Electrophoresis
10 18:409-417 (1997)). The resulting peptides were
extracted and the mass of approximately 30 peptides was
determined using a Bruker Reflex MALDI-TOF mass
spectrometer, providing a unique signature by which to
identify the protein by peptide mass searches (Mann et
15 al, Biol. Mass Spectrom. 22:338-345 (1993), Pappin et al,
J. Exp. Med. 180:273-281 (1994)). The 55 kDa angiostatin
binding membrane protein was identified as the α/β
subunits of ATP synthase (Table I), whereas the
plasminogen binding protein was confirmed as annexin II.
20 Although expression of the β subunit of ATP synthase has
been reported on the surface of several tumor cell lines
(Das et al, J. Exp. Med. 180:273 (1994)), this is the
first evidence for surface expression of the subunits of
ATP synthase on HUVEC.

Table I

Bruker Reflex MALDI-TOF mass spectrometer
analysis of 55 kDa peptides

5

Sequence	peptide mass (monoisotopic)	
	Measured (Da)	Calculated (Da)
QMSLLLR	859.48	859.495
AVDSLVPIGR	1025.58	1025.587
VGLKAPGIIPR	1119.68	1119.713
TIAMDGTEGLVR	1261.40	1261.634
ISVREPMQTGIK	1357.70	1357.739
IMNVIGEPIDER	1384.68	1384.702
AHGGYSVFAGVGER	1405.66	1405.674
FTQAGSEVSALLGR	1434.73	1434.747
TSIADTIINQKR	1471.81	1471.836
EAYPGDVFYLHSR	1552.71	1552.731
VALVYGQMNEPPGAR	1600.79	1600.803
TGAIVDVPVGEELLGR	1623.87	1623.883
LVLEVAQHLGESTVR	1649.88	1649.910
IMDPNIVGSEHYDVAR	1814.85	1814.862
VLDSGAPIKIPVGPETLGR	1918.08	1918.089
AIAELGIYPAVDPLDSTSR	1986.99	1987.026
IMNVIGEPIDERGPICK	2009.10	2009.098
IPSAVGYQPTLATDMGMTQER	2265.06	2265.077
EVAAFAQFGSDLDAATQQLSR	2337.15	2337.160

EXAMPLE 4

Binding of the α Subunit ATP Synthase
Antibody to the Surface of HUVEC by
Flow Cytometry and Immunofluorescence Microscopy

5 To further confirm the surface localization of the
ATP synthase, HUVEC were analyzed by flow cytometry and
immunofluorescence microscopy. A rabbit polyclonal
antiserum raised against the α subunit of ATP synthase
from *E. coli* reacted with the cell membranes of HUVEC as
10 determined by fluorescence-activated flow cytometry
(Fig. 4). Control flow cytometry studies were performed
using A549 cells which are known to express the α/β
subunits of ATP synthase (Das et al, J. Exp. Med.
180:273-281 (1994)) (Fig. 4B). A549 cells were also
15 analyzed with anti- α subunit ATP synthase antibody pre-
incubated with a 5-fold molar excess of recombinant α
subunit of ATP synthase protein and showed a decreased
affinity for binding (Fig. 4C). HUVEC were incubated
with increasing concentrations of antibody to determine
20 saturation. Fig. 4D demonstrates specific, saturable
binding of antibody directed against the α subunit of ATP
synthase on HUVEC membranes.

Immunofluorescence microscopy of HUVEC confirmed
the surface-associated immunoreactivity of α subunit
25 ATP synthase antibody on HUVEC cell membranes
(Fig. 5A). Control experiments were performed with
secondary antibody alone (Fig. 5D), pre-immune serum
(Fig. 5E) and permeabilized HUVEC in the presence of
anti- α subunit ATP synthase antibody (Fig. 5F). Human

dermal microvascular endothelial cells also reacted with antiserum raised against the α subunit of ATP synthase (Fig. 5C).

5

EXAMPLE 5

Inhibition of Angiostatin Binding in the Presence of the Antibody to the α Subunit of ATP Synthase

The rabbit polyclonal antiserum raised against the α subunit ATP synthase blocked binding of angiostatin to HUVEC by 59%, demonstrating that this protein functions in angiostatin binding (Fig. 6). In addition, ^{125}I -labeled angiostatin bound to purified recombinant α subunit from human ATP synthase (Fig. 7B) and binding was inhibited ~56% in the presence of a 250-fold molar excess of unlabeled angiostatin (Fig. 7C). Complete inhibition of binding was not obtained and may be due in part to non-specific binding, improper folding of the recombinant protein or binding epitopes only found in the presence of the α/β heterodimer. Furthermore, binding to the recombinant α subunit ATP synthase was not inhibited by a 2500-fold molar excess of unlabeled plasminogen (Figure 7D). Further, ^{125}I -labeled plasminogen did not bind to the recombinant α subunit ATP synthase (Figure 7E), but did bind to annexin II (Figure 3D).

EXAMPLE 6

Inhibition of Proliferation in the Presence of Antibody
to the α Subunit of Human ATP Synthase

To determine whether the anti-proliferative
5 effects of angiostatin were mediated by ATP synthase
binding, cell proliferation assays were performed in
the presence of antiserum raised against the α subunit
of ATP synthase from *E. coli*. The inhibitory effects
of angiostatin on HUVEC proliferation were abrogated by
10 ~81% in the presence of antibody to the α subunit of
ATP synthase (Table II), providing direct evidence that
angiostatin binding to the α subunit of ATP synthase
functions as a mechanism for inhibition of endothelial
cell growth. These data indicated that this binding
15 site serves as a receptor for angiostatin.

Table II The anti-proliferative effect of angiostatin is reversed by anti- α subunit ATP synthase antibody.

Concentration angiostatin added, μ M	Percent proliferation inhibited, +/-SEM		
	Without antibody	With antibody	% Recovery
0	0	0	0
0.5	10 \pm 1.4	1 \pm 0.2	90
0.75	25 \pm 4.2	5 \pm 4.1	80
1.0	23 \pm 9.0	6 \pm 0.8	74

5 HUVEC were plated at a density of 5000 cells/well in media containing angiostatin at a final concentration of 0.5, 0.75 and 1.0 μ M. Anti- α subunit ATP synthase antibody derived from *E. coli* was added concomitantly at a dilution of 1:10. MTS/PMS solution was added and
10 absorbance of formazan was quantitated according to the manufacturer's specifications (Promega, Madison, WI). The average proliferative effect of pre-immune serum and anti- α subunit antibody alone increased 4.6% over
15 buffer control. Results represent three separate experiments performed in duplicate with S.E.M. Percent recovery represents the ability of the anti- α subunit ATP synthase antibody to block the antiproliferative effect of angiostatin, and thereby restore
20 proliferation to an average of 81% of that obtained with the control cells.

ATP synthase is composed of two functional domains termed F_1 and F_0 . The F_1 portion contains multiple
25 subunits ($\alpha_3\beta_3\gamma\delta\epsilon$) and acts as the catalytic site for ATP synthesis, while the membrane embedded F_0 portion is a proton channel (Penefsky et al, Advances in

Enzymology and Related Areas of Molecular Biology
64:173-214 (1991)). Isolated α and β subunits bind ATP
and have weak ATPase activity; however, ATP synthesis
requires all F_1 and F_0 subunits (Boyer, Ann. Rev.
5 Biochem, 66:717-749 (1997)).

Endothelial cells play a strategic role within the
vasculature, serving as a barrier between the
intravascular compartment and the underlying tissues and
are often exposed to hypoxic stress. Relative to other
10 cell types, endothelial cells are more resistant to
hypoxic challenge by their ability to maintain a high
level of intracellular ATP (Graven et al, Kidney
International 51:426-437 (1997)). A plasma membrane-
associated ATP synthase may produce extracellular ATP
15 which can diffuse back into the cell providing an
additional, albeit limited, ATP source (Unno et al, Am.
J. Physiol. 270:g1010 (1996), Unno et al, Surgery 121:668
(1997)). Angiostatin, by binding to the α/β subunits of
plasma membrane-localized ATP synthase, may disrupt this
20 production of ATP, rendering endothelial cells more
vulnerable to hypoxic challenge and eventual irreversible
cell damage. In the microenvironment of a growing tumor,
tissue hypoxia provides a powerful stimulus for the
production of angiogenic growth factors such as VEGF,
25 bFGF, and angiopoetin. The ability of host endothelial
cells to respond to these growth factors by increased
proliferation is likely dependent on the ability to
survive hypoxic challenge. By abolishing the ability to
resist low oxygen tension, angiostatin may decrease

endothelial cell survival in the tumor microenvironment. It has recently been reported that angiostatin may also function by inducing endothelial cell apoptosis, providing an additional independent mechanism for the anti-angiogenic action of this polypeptide (Claesson-Welsh et al, Proc. Natl. Acad. Sci., 95:5579-5583 (1998)).

EXAMPLE 7

10 Binding of Antibody Directed Against the β Subunit of ATP Synthase on the Surface of HUVEC by Flow Cytometry

HUVEC cells were resuspended in ice-cold staining buffer (HBSS, 1% BSA, 0.1% sodium azide) and incubated on ice for 30 min with either rabbit polyclonal anti-serum raised against β subunit ATP synthase derived from *E. coli* or pre-immune rabbit serum. Cells were washed with ice-cold staining buffer and pelleted in a microfuge at 4°C. This wash was repeated twice and the cells resuspended in ice-cold staining buffer prior to incubation on ice for 30 min in the dark with goat anti-rabbit IgG conjugated to fluorescein isothiocyanate (FITC). Following the final wash (as above), the cells were pelleted and fixed in 10% neutral buffered formalin at a density of 1×10^6 cells/ml. Control experiments were performed using antibody directed against the β subunit of ATP synthase which was preincubated with a 5-fold molar excess of recombinant β subunit ATP synthase protein. The mean relative fluorescence following

excitation at a wavelength of 488nm was determined for each sample on a FACScan flow cytometer (Becton-Dickenson) and analyzed with CellQuest software (Becton-Dickenson).

5

* * *

All documents cited above are hereby incorporated in their entirety by reference.

One skilled in the art will appreciate from a reading of this disclosure that various changes in form and detail can be made without departing from the true scope of the invention.

10

WHAT IS CLAIMED IS:

1. A method of screening a test compound for its ability to inhibit or enhance the binding of angiotensin to ATP synthase comprising:

i) contacting said test compound and angiotensin with ATP synthase, or angiotensin binding portion thereof, under conditions such that angiotensin can bind to said ATP synthase, or angiotensin binding portion thereof, in the absence of said test compound, and

ii) determining the amount of angiotensin bound to said ATP synthase, or angiotensin binding portion thereof, and comparing that amount to an amount of angiotensin bound to said ATP synthase, or angiotensin binding portion thereof, in the absence of said test compound,

wherein a reduction in the amount of angiotensin bound to said ATP synthase, or angiotensin binding portion thereof, in the presence of said test compound indicates that said test compound inhibits the binding of angiotensin to said ATP synthase, or angiotensin binding portion thereof, and

wherein an increase of the amount of angiotensin bound to said ATP synthase, or angiotensin binding portion thereof, in the presence of said test compound indicates that said test compound enhances the binding of angiotensin to said ATP synthase, or angiotensin binding portion thereof.

2. The method of claim 1 wherein said angiostatin bears a detectable label.
3. The method of claim 1 wherein said ATP synthase, or angiostatin binding portion thereof, is attached to a solid support.
4. The method of claim 1 wherein said ATP synthase, or angiostatin binding portion thereof, is associated with a lipid membrane.
5. The method of claim 4 wherein said membrane is a membrane of an intact cell.
6. The method of claim 5 wherein said cell naturally expresses ATP synthase.
7. The method of claim 5 wherein said cell has been transformed with a nucleic acid sequence that encodes said ATP synthase, or angiostatin binding portion thereof.
8. A compound identified in the method of claim 1 as inhibiting the binding of angiostatin to said ATP synthase or angiostatin binding portion thereof.
9. A compound identified in the method of claim 1 as enhancing the binding of angiostatin to said ATP synthase or angiostatin binding portion thereof.
10. A method of screening a test compound for its ability to modulate a bioactivity resulting from binding of angiostatin to ATP synthase comprising:

i) contacting said test compound and angiostatin with a cell that expresses ATP synthase, or angiostatin binding portion thereof, under conditions such that angiostatin can bind to said ATP synthase, or angiostatin binding portion thereof, in the absence of said test compound, and

ii) determining the amount of angiostatin required to achieve the same bioactivity in the presence of said test compound as in the absence of said test compound,

wherein a reduction in the amount of angiostatin required to achieve said same bioactivity in the presence of said test compound indicates that said test compound is an angiostatin agonist, and

wherein an increase in the amount of angiostatin required to achieve said same bioactivity in the presence of said test compound indicates that said test compound is an angiostatin antagonist.

11. An angiostatin agonist identified in accordance with the method of claim 10.

12. An angiostatin antagonist identified in accordance with the method of claim 10.

13. The method of claim 10 wherein said bioactivity is inhibition of cell proliferation.

14. The method of claim 10 wherein said bioactivity is enhancement of proton pumping.

15. A method of inhibiting the angiogenesis inhibitory effect of angiostatin in a patient comprising

administering to said patient an amount of an angiostatin antagonist that binds an angiostatin binding portion of ATP synthase sufficient to effect said inhibition.

16. A method of inhibiting the angiogenesis inhibitory effect of angiostatin in a patient comprising administering to said patient an amount of a soluble angiostatin binding portion of ATP synthase sufficient to effect said inhibition.

17. A method of enhancing the angiogenesis inhibitory effect of angiostatin in a patient comprising administering to said patient an amount of an angiostatin agonist that binds to an angiostatin binding portion of ATP synthase sufficient to effect said enhancement.

18. An expression construct comprising a vector and a nucleic acid sequence encoding the α subunit of ATP synthase, or angiostatin binding portion thereof, operably linked to a promoter.

19. A host cell comprising the construct of claim 18.

20. A method of producing the α subunit of ATP synthase, or angiostatin binding portion thereof, comprising culturing the host cell of claim 19 under conditions such that said nucleic acid is expressed and said α subunit of ATP synthase, or angiostatin binding portion thereof, is thereby produced.

21. An antibody specific for the α subunit of ATP synthase, or angiostatin binding portion thereof, or antigen binding portion thereof.

22. A kit comprising ATP synthase, or angiostatin binding portion thereof, and angiostatin, or truncated form thereof.

23. An isolated complex comprising angiostatin and ATP synthase, or angiostatin binding portion thereof.

24. The complex according to claim 23 wherein said complex is bound to a solid support.

1/9




Fig.1A




Fig.1B

2/9




Fig.2




Fig.3

4/9




Fig. 4A




Fig. 4B





Fig.4D

Fig.5C



Fig.5B




Fig.5A




Fig.5F




Fig.5E




Fig.5D



7/9




Fig.6




Fig.7

9/9




Fig.8

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/10957

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

DIALOG

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BOYER, P.D. The ATP Synthase--A Splendid Molecular Machine. Annual Review Biochemistry. 1997, Vol. 66, pages 717-749.	1-24
X	KATAOKA et al. Nucleotide Sequence of a cDNA for the Alpha Subunit of Human Mitochondrial ATP Synthase. Biochimica et Biophysica Acta. 1991, Vol. 1089, pages 393-395, see entire document.	18-20
X	RAO et al. The Defective Proton-ATPase of uncA Mutants of Escherichia coli: ATP-Binding and ATP-Induced Conformational Change in Mutant Alpha-Subunits. June 1987, Vol. 255, No. 2, pages 309-315, especially page 310, column 2.	8, 12, 21

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

14 SEPTEMBER 1999

Date of mailing of the international search report

14 OCT 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer:

JAMES L. GRUN, PH.D.

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/10957

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DAS et al. A Novel Ligand in Lymphocyte-Mediated Cytotoxicity: Expression of the Beta Subunit of H ⁺ Transporting ATP Synthase on the Surface of Tumor Cell Lines. Journal of Experimental Medicine. July 1994, Vol. 180, pages 273-281.	1-17
X	US, A, 5,639,725 (O'REILLY et al.) 17 June 1997, see entire document, especially columns 6-7, 10-11, 17-18.	8, 11, 12, 17, 22-24
----		-----
A		1-7, 9-10, 13-16

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/10957

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A61K 38/17, 38/43; C07H 21/04; C07K 14/435, 14/705, 16/28, 16/40; G01N 33/53, 33/543, 33/554

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

424/143.1, 152.1, 155.1, 172.1; 435/7.21, 69.1, 70.1, 320.1; 436/503, 518, 519; 514/12; 530/350, 380; 536/23.5; 935/22, 66

B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

424/143.1, 152.1, 155.1, 172.1; 435/7.21, 7.23, 7.8, 7.92, 13, 69.1, 70.1, 320.1, 330, 334, 337, 344, 375; 436/503, 518, 519; 514/12; 530/350, 380, 388.22, 388.25, 388.8, 389.3; 536/23.5; 935/22, 66

